



Fungi-mix

Indication:

Fungal infections such as Candidiasis, Moniliasis and Aspergillosis in birds

Yeast are found everywhere in the environment including spoiled food. For this reason, fresh fruits and vegetables should not be left in the bird's cage for long periods of time. In the summer heat and humidity, food can spoil in as little as a few hours. Poor hygiene can also spread yeast infections. Not sterilizing hand feeding equipment, improper cage cleaning and even not washing our hands can spread the yeast. Most healthy birds will not be affected. Older birds, neonates, sick birds and stressed birds generally develop yeast infections as secondary infections. These secondary infections put increased stress on their already overtaxed immune system and interfere with the healing process.

Symptoms:

Flung out of the feathers • Bad appetite • Vomiting • Swollen crop
Breathing difficulties • Often combined with diarrhoea.

Dosage

Dose eggfood: birds in general: 5 g / 500 g of soft food, twice a day, 5 to 7 days

Dose drinkingwater: birds in general: 5 g/1 L drinkingwater, twice a day, 5 to 7 days. (stir well before use!) Nystatin can be mixed directly into the handfeeding formula but is more effective if given full strength about 1/2 hour before feeding. This will provide it time to coat the crop lining and to attack the unhealthy Candida organism. For prevention provide your birds citric acid 5 ml per 1 liter drinkingwater daily. Measuring spoon 1+5 g. included.



Interaction(s): not known. **Contra-indication(s):** over dosage.

Advice for use:

Isolate the sick birds (temperature 28-33 °C, lights dimmed).

While treating sick animals, provide preventive treatment to other birds as well.

Ensure that the birds take the recommended quantities of medication; therefore, avoid the intake of non medicated food or water.

Administer glucose and MULTIVIT MIX during treatment.

For serious conditions, extended treatment is recommended

Make sure to adequately disinfect the accommodation.

If rapid improvement fails to occur, consult an expert.

NOTES: READ INSTRUCTIONS BEFORE USE.

Store dry, out of reach of direct sunlight, between 0-30 °C

Shelf life after first opening; One (1) year.

Batchnr. and expiry date: see package.

Avoid contact of the medicine with your hands.

Keep out of reach of children.

For animal treatment only. Not for animals intended to produce food for human consumption

Dispose of empty packaging and remaining product in the household refuse.

This veterinary medicine is marketed in accordance with the small animal exemption scheme

Contains:

100 g Nystatine (200.000 i.u./gram)