



Orni-mix

Indication:

Infections of the airways caused by **Ornithosis, Pneumonia, Mycoplasmosis, Psittacosis (Chlamydiosis)** in birds.

Symptoms:

- Sinus infection, labored breathing, shortness of breath, noisy breathing or 'runny nose'
- Eye discharge or swelling, matting of feathers around eyes
- Lethargy, rued / fluffed-up feathers, loss of appetite, weightloss, weakness.

Dosage

Dosage: Cage-and aviary birds(canaries, finches):

5 g / 2 liters of drinkingwater, 5-10 consecutive days.

Dosage: Parakeets, Parrots:

1 g/2 kg of body weight, or 3 g/200 ml drinkingwater, 5-10 consecutive days.

For treatment of Psittaccosis (Parrot fever): 1 g/2 kg of body weight, or 3 g/200 ml drinkingwater. Duration of treatment is for 45 days.

All sources of calcium need to be removed for this period, as calcium aects the uptake of the antibiotic.

Measuring spoon 1+5 g. included.

Interaction(s): not known. **Contra-indication(s):** over dosage.

Advice for use:

Isolate the sick birds (temperature 28-33 °C, lights dimmed).

While treating sick animals, provide preventive treatment to other birds as well.

Ensure that the birds take the recommended quantities of medication; therefore, avoid the intake of non medicated food or water.

Administer glucose and MULTIVIT MIX during treatment.

For serious conditions, extended treatment is recommended

Make sure to adequately disinfect the accommodation.

If rapid improvement fails to occur, consult an expert.

NOTES: READ INSTRUCTIONS BEFORE USE.

Store dry, out of reach of direct sunlight, between 0-30 °C

Shelf life after rst opening; One (1) year.

Batchnr.and expiry date: see package.

Avoid contact of the medicine with your hands.

Keep out of reach of children.

For animal treatment only. Not for animals intended to produce food for human consumption

Dispose of empty packaging and remaining product in the household refuse.

This veterinary medicine is marketed in accordance with the small animal exemption schem

Contains:

Doxycyline Hyclate 6%, Spiramycin Adipate 10% Metronidazole 10%, Bromhexine HCl 2%, inactive ingredients q.s to 100 g

