



Salmo-mix

Indication:

Salmonellosis, colibacillosis (sweating disease) and bacterial infections common in birds.

Symptoms:

SYMPTOMS OF BACTERIAL INFECTIONS IN COMMON:

• Obvious changes in behaviour and appearance • ruffled and soiled feathers • Diarrhoea • Unusual or difficult breathing • reduced appetite and sitting low on perch

EXTRA SYMPTOMS OF COLIBACILLOSIS (SWEATING DISEASE):

• Inside of nests stained with yellow diarrhoea of nestling • youngster stunted • increased death rate during the first few days of life.

Salmonellosis is more prevalent in wetter periods of the year, outside aviary, when proper hygienic measures are not taken, exposure to mice or wild bird droppings is a source of infection.



Dosage

DOSAGE: AS A NORMAL TREATMENT add a dosage of 5 g in 1 liter of drinkingwater during 5 –7 successive days.

IN CASE OF SALMONELLOSIS OR COLIBACILLOSIS (SWEATING DISEASE) repeat the treatment after 2 days

Measuring spoon 1+5 g. included.



Interaction(s): not known. **Contra-indication(s):** over dosage.

Advice for use:

Isolate the sick birds (temperature 28-33 °C, lights dimmed).

While treating sick animals, provide preventive treatment to other birds as well.

Ensure that the birds take the recommended quantities of medication; therefore, avoid the intake of non medicated food or water.

Administer glucose and MULTIVIT MIX during treatment.

For serious conditions, extended treatment is recommended

Make sure to adequately disinfect the accommodation.

If rapid improvement fails to occur, consult an expert.

NOTES: READ INSTRUCTIONS BEFORE USE.

Store dry, out of reach of direct sunlight, between 0-30 °C

Shelf life after first opening; One (1) year.

Batch number and expiry date: see package.

Avoid contact of the medicine with your hands.

Keep out of reach of children.

For animal treatment only. Not for animals intended to produce food for human consumption

Dispose of empty packaging and remaining product in the household refuse.

This veterinary medicine is marketed in accordance with the small animal exemption scheme

Contains:

Furaltadone HCl 12,8%, Chloramphenicol 8%, Chlortetracycline HCl 12,5%, Inactive ingredients q.s to 100 g