



Worm-mix

Indication:

For prevention and treatment of round-and hairworm infestations in birds.

Symptoms:

OF INTESTINAL WORMS

- Weight loss
- Loss of condition
- Increased vulnerability to other diseases.

WHEN DO YOU TREAT YOUR ANIMALS:

- As a routine, after moult
- If you buy new birds
- In case of worm infections

Dosage

DOSAGE: AS A GENERAL TREATMENT FOR BIRDS:

1 g for 500 g live-weight /5 g for 2500 g live-weight for 2 consecutive days

Administer the Worm-mix with the feed or in the drinking water.

Repeat this treatment after 10-12 days.

Measuring spoon 1+5 g. included.

Interaction(s): not known. **Contra-indication(s):** over dosage: Don't administer this product during coupling, feeding of youngsters, moulting of big feathers or to birds younger than 14 days old.

Advice for use:

Isolate the sick birds (temperature 28-33 °C, lights dimmed).

While treating sick animals, provide preventive treatment to other birds as well.

Ensure that the birds take the recommended quantities of medication; therefore, avoid the intake of non medicated food or water.

Administer glucose and MULTIVIT MIX during treatment.

For serious conditions, extended treatment is recommended

Make sure to adequately disinfect the accommodation.

If rapid improvement fails to occur, consult an expert.

NOTES: READ INSTRUCTIONS BEFORE USE.

Store dry, out of reach of direct sunlight, between 0-30 °C

Shelf life after first opening; One (1) year.

Batch number and expiry date: see package.

Avoid contact of the medicine with your hands.

Keep out of reach of children.

For animal treatment only. Not for animals intended to produce food for human consumption

Dispose of empty packaging and remaining product in the household refuse.

This veterinary medicine is marketed in accordance with the small animal exemption scheme

Contains:

Levamisole HCl 2%, Inactive ingredients q.s. to 100 g

