



Orni Special

Indication:

Ornithosis, Mycoplasmosis, Respiratory problems and Canker.

Symptoms:

Picking of feathers, scratching head, red throat, yawning, inflamed third eyelid (especially in young pigeons), dirty noses, blue flesh, swollen heads, one eye cold, wet eyes, snot.

Dosage

Prevention: 1 measuring spoon (=5g) for 2 litres drinking water during 1 or 2 days every 2 weeks, preferably starting the day after the race.

As a treatment: 1 measuring spoon (=5g) for 2 litres drinking water during 5-7 successive days.

This cure may be repeated if necessary after a few days.

Stir well before use!

Orni Special is a water-soluble powder which can also be given under the feed.

Measuring spoon 1+5 g included.

Advice for use:

- Put the sick animals in quarantine.
- Treat all of your birds, not only the sick ones.
- Mix the needed dose in an amount of drinking water or feed the animals consume within one day.
- Be sure no other water is available during the treatment.
- Do not provide any calcium, grit, salt or dairy products during treatment.
- After medication give your animals MULTIVITAMIN (the day after last day of treatment).
- Give a maintenance dose of MULTIVITAMIN every week
- Disinfect lofts/houses.
- Do not give the day of shipping.
- In case of doubt or persistent problems, consult a vet. surgeon.

NOTES: READ INSTRUCTIONS BEFORE USE.

- Store dry, out of reach of direct sunlight, between 0-30 °C.
- Shelf life after first opening; One (1) year.
- Batchnr. and expiry date: see package.
- Avoid contact of the medicine with your hands.
- Keep out of reach of children
- Interaction(s): not known. Contra indication(s) administration with Furaladone, and overdosage.
- For animal treatment only. Not for animals intended to produce food for human consumption.
- Dispose of empty packaging and remaining product in the household refuse.
- This veterinary medicine is marketed in accordance with the small animal exemption scheme.

Contains:

Doxycyline Hyclate 6%, Spiramycin Adipate 10%, Metronidazole 10%, Bromhexine HCl 2%, Inactive ingredients q.s. to 100 g.

