



Worm-Stop

Indication:

Indication: Hair -and roundworm infestations in pigeons
(*Capilarias spp* and *Ascaridia spp*)

Symptoms:

The birds lose weight. Heavily infested pigeons usually show impaired general condition and are listless and apathetic with dull, ruffled plumage and droppings of varying consistency.



Dosage

5 g. for 2 L. drinking water, (40 pigeons), 1 or 2 days, Pigeons should not be dewormed while they are rearing the young or during the main moult, repeat the treatment after 10-12 days.

Worm-Stop is a water-soluble powder which can also be given under the feed.

Measuring spoon 1+5 g included



Advice for use:

- Put the sick animals in quarantine.
- Treat all of your birds, not only the sick ones.
- Mix the needed dose in an amount of drinking water or feed the animals consume within one day.
- Be sure no other water is available during the treatment.
- After medication give your animals MULTIVITAMIN (the day after last day of treatment).
- Give a maintenance dose of MULTIVITAMIN every week.
- Disinfect lofts/houses.
- Do not give the day of shipping.
- In case of doubt or persistent problems, consult a vet. surgeon.

NOTES: READ INSTRUCTIONS BEFORE USE.

- Store dry, out of reach of direct sunlight, between 0-30 °C.
- Shelf life after first opening; One (1) year.
- Batchnr. and expiry date: see package.
- Avoid contact of the medicine with your hands.
- Keep out of reach of children.
- Interaction(s): not known. Contra indication(s) overdose.
- For animal treatment only. Not for animals intended to produce food for human consumption.
- Dispose of empty packaging and remaining product in the household refuse.
- This veterinary medicine is marketed in accordance with the small animal exemption scheme.

Contains:

Levamisole HCl 16% , Inactive ingredients q.s. to 100 g.

